

**“Combination therapy for reduction of toxicity of chemotherapeutic agents”**

5 The present invention relates to a unique combination therapy for minimising dosage required of chemotherapeutic agents for their use in antiviral, antibacterial, antiparasitic and anticancer chemotherapy. This invention also relates to the use of the compounds of the present invention including precursor molecules, derivatives, metabolites, analogues, mimic molecules and to compositions containing the compounds of the present invention including precursor molecules, derivatives, metabolites, analogues, mimic molecules for use in a combination  
10 therapy with chemotherapeutic agents for their use in antiviral, antibacterial, antiparasitic and anticancer chemotherapy.

Cancer develops from changes in the DNA, or genetic material, of the body's cells, causing them to develop into precancerous lesions. Such lesions exhibit a strong  
15 tendency to develop into malignant tumours, or cancer. Such lesions include lesions of the breast (that can develop into breast cancer), lesions of the skin (that can develop into malignant melanoma or basal cell carcinoma), colonic adenomatous polyps (that can develop into colon cancer), and other such neoplasms.

20 Cancer may take years to develop. The process typically begins with some disruption to the DNA of a cell, the genetic code that directs the life of the cell. Many things, such as diet, tobacco, sun exposure or certain chemicals can cause such disruptions. Some cells will enter a precancerous phase, known as dysplasia. Some cells will also enter the state of *carcinoma in situ*, in which the cancer cells  
25 are restricted to a microscopic site and do not pose a great threat. Eventually, unless the body's own immune system takes care of the wayward cells either on its own or by being enhanced by specific chemicals, a tumour will develop. It may take as long as 30 years for a tumour to go through the entire process and become large enough to produce clinical symptoms.

30 Anyone can get cancer, including children, but it is most common in people over the age of 50. This year about 1.22 million people in the United States will be

diagnosed with cancer (not including the more than 1 million annual cases of basal and squamous-cell skin cancers.) About 563,000 people will die of cancer this year.

Treatment for cancer has progressed rapidly over the last 30 years. Doctors generally prescribe three main treatments for cancer: surgery, radiation therapy, chemotherapy or a combination of these. Choosing a course of medical treatment depends largely on the cancer type, stage of progression, and location.

Chemotherapy uses poison drugs that take advantage of cancer cells' rapid growth and consumption of large amounts of nutrients. Chemotherapy side effects include nausea, bone marrow suppression and temporary full or partial hair loss. One class of chemotherapeutic agents, the nucleotide analogues, are used as anticancer agents because they interfere with the synthesis of DNA and thereby preferentially kill rapidly dividing cells such as tumour cells. Some of the nucleotide analogues commonly used in chemotherapy are 6-mercaptopurine, 5-flourouracil, 5-iodo-2'-deoxyuridine and 6-thioguanidine. Each of these compounds disrupts the normal replication process by interfering with the formation of correct Watson-Crick base-pairing.

Physicians inject these drugs into the bloodstream, where they travel throughout the body, consumed by every cell. Rapidly growing cancerous cells consume much more of the poisonous drugs than do normal cells. As a result, the drugs destroy cancerous cells faster than normal cells. However, chemotherapy drugs act on all the patient's cells -- the cancerous cells and the healthy cells. A physician's challenge is to administer the drugs to kill only the cancer cells, not the healthy cells. Unfortunately, most chemotherapeutic drugs have serious side effects that prohibit their long-term use, or use in otherwise healthy individuals with precancerous lesions. These side effects, which are a result of non-specific toxicity of the drugs, immunosuppression and other toxicities. For this reason there is a need to identify new drug candidates for therapy of patients with precancerous lesions that can be combined with existing chemotherapeutic drugs, whereby the dose can be reduced, diminishing the toxicity and serious side effects in humans.

Acquired Immunodeficiency Syndrome (AIDS) is one of the most significant infections to appear in the last decade. This epidemic is not confined to a single segment of the population nor is its spread blocked by natural barriers or

international boundaries. Millions have died in Africa and many more individuals are infected worldwide. In the United States more than 100,000 people have died and at least 1 million more are presently infected with the virus. This pandemic shows no signs of abating.

5

AIDS was first diagnosed in male homosexuals who exhibited a variety of infections of fungal (*Candida albicans*), protozoal (*Pneumocystis carinii*), and viral (Herpes zoster) origin. Many of these individuals also had an increased incidence of kaposi sarcoma and lymphoma. They had a depressed T helper/T suppressor lymphocyte cell ratio and an absence of delayed hypersensitivity responses. Collectively, these observations suggested a deficiency in cell-mediated immunity.

10

It is strongly suspected that the causative agent in AIDS is an RNA retrovirus called the human immunodeficiency virus (HIV-1 or HIV-2). HIV possesses an envelope glycoprotein (gp120) that has a high affinity for the CD<sub>4</sub> receptor on T helper cells and other target cells. These other target cells include bone marrow stem cells, macrophages, endothelial cells, glial cells, lymph node, dendritic cells, bowel enterochromaffin cells, cervical epithelium and possibly Langerhans cells. However, it is the effects of HIV on T-helper cells that are the best known. The infectious process begins when the virus penetrates the body and enters the blood stream. Binding of HIV to CD<sub>4</sub> target cells involves interaction of the external envelope glycoprotein molecule gp120 with the CD<sub>4</sub> molecule, although other cell receptors may be involved. The virus next enters the target cell, or is internalized, through fusion of the viral envelope with the target cell membrane. Through this fusion, the virus loses its coat, and releases its RNA core and reverse transcriptase enzyme into the host cell cytoplasm.

15

20

25

The HIV reverse transcriptase enzyme copies the RNA message producing first a single-stranded, and then a double-stranded, DNA (circular complementary DNA). This newly formed double-stranded DNA becomes incorporated into the host chromosomal DNA once it enters the host cell nucleus. This incorporated viral DNA may remain dormant or, upon activation, will produce viral messenger RNA (mRNA). The viral mRNA codes for proteins that are important in viral replication.

30

Glycoprotein will then envelop the RNA genome resulting in the production of infectious viral particles; completed viral particles are then released to infect other cells.

5 Current approaches to HIV treatment generally involve immunotherapy (e.g., vaccines against whole killed HIV and a variety of HIV surface glycoproteins) directed at the HIV as well as pharmacological intervention in the HIV infectious process. In theory, any of the steps of viral replication or release could be points of pharmacological attack against the virus. The major chemotherapeutic attack by  
10 available drugs has been at the level of inhibition of viral reverse transcriptase using nucleotide and nucleotide analogues. The first drug licensed for use in HIV treatment became available in 1987; it was azidothymidine (AZT). In the early 1990's, dideoxyinosine (DDI) and dideoxycytidine (DDC) were approved by the FDA. AZT and DDI were approved for monotherapy while DDC is used in  
15 combination with one of the other drugs. However, when administered to patients, nucleotide and nucleoside analogues have shown toxicity to liver, bone marrow, and the nervous system. Side effects include bone marrow suppression (anemia granulocytopenia), headache, malaise, nausea, pancreatitis, peripheral neuropathy, minor GI and CNS symptoms, and oral ulcers. In the case of antiviral therapy,  
20 nucleoside analogs have been rarely curative, and the side effects that arise during chronic administration of the drug often cause therapy to be discontinued or altered. In the case of cancer therapy, where intent is to kill the cancer cells, the compound and protocol (dose, method of administration, timing of doses) must be carefully designed and monitored to minimize the damage to non-cancerous  
25 tissues.

Many different treatment regimens are and have been used to treat the HIV infection and AIDS, which occurs after the latent infection. While they might prolong survival and possibly minimize symptoms, in view of the mounting worldwide  
30 concern regarding the epidemic, these treatments have not been generally successful and cause considerable side effects. Therefore, the continuing hard reality is that once the virus enters the body and begins the uncoating process, a fatal outcome is almost inevitable. Such an outcome reveals the continuing need

for additional research to discover a method of treatment, which can suppress the reproduction of latent viruses such as HIV without inducing toxicity. For this reason there is a need to identify new drug candidates for therapy of patients with HIV infection and AIDS that can be combined with existing chemotherapeutic  
5 drugs, whereby the dose can be reduced, diminishing the toxicity and serious side effects in humans.

Parasitic infections are a major worldwide health problem. The global prevalence of human parasitic infections already exceeds 50% and is increasing. Diverse factors  
10 are responsible including population crowding; poor sanitation and health education; inadequate control of parasite infections and reservoirs of infection; increased world travel; population migration and resistance to the agents used for chemotherapy or control of vectors.

15 Despite encouraging progress in identifying promising molecular targets for intervention with vaccines, chemotherapy remains the single most effective method to control most parasitic infections. However, to be used for mass chemotherapy, an ideal antiparasitic agent should be safe at high therapeutic doses with minimal toxicity. Few antiparasitic drugs meet these criteria. For this reason there is a  
20 need to identify new drug candidates for therapy of patients with parasitic infections that can be combined with existing chemotherapeutic drugs, whereby the dose can be reduced, diminishing the toxicity and serious side effects in humans. Furthermore, several antibiotics are not in current use due to their toxicity profile. It would be advantageous similarly to identify new drug candidates for therapy of  
25 patients with bacterial infections that can be combined with existing chemotherapeutic drugs, whereby the dose can be reduced, diminishing the toxicity.

30 Additionally, the suppression of immune response to limit rejection in tissue transplantation by the use of nucleotide and nucleoside analogues is limited by the toxicity of the compounds. The identification of new drug candidates for therapy of patients with tissue transplants that can be combined with existing chemotherapeutic drugs, whereby the dose can be reduced, is necessary.

5 The correlation between the combination therapy including the compounds of the present invention, circiliol, precursor molecules, derivatives, metabolites, analogues and/or mimic molecules, antiviral and anticancer chemotherapy was not recognised prior to the work of the applicant. Accordingly the following provides information on each of these topics.

10 The present invention provides methods and pharmaceutical compositions for repressing reproduction of latent viruses, such as HIV, parasites, bacteria and for anticancer therapy in humans and animals, by the generally concurrent administration of at least one chemotherapeutic agent with compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof.

15 The present invention relates to a composition comprising at least one of the compounds of the present invention.

20 The present invention also relates to a pharmaceutical formula comprising a composition comprising at least one of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof, at least one chemotherapeutic agent and a pharmaceutically acceptable carrier. The advantage of such a pharmaceutical formula is that the antimetabolites, nucleotide and/or nucleoside analogues can be administered in much lower doses than normally administered, thereby greatly  
25 reducing their toxicity while increasing their therapeutic index. The synergistic effect of the chemotherapeutic agents in combination with the compounds of the present invention provides an enhanced efficacy despite the low dose of chemotherapeutic agent administered.

30 The present invention is also directed to the use of a pharmaceutical formulation for the manufacture of a medicament for treatment of a mammal suffering from a neoplasia and/or viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation.

The present invention is also directed to a method for manufacturing a medicament for treating neoplasia and/or viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation utilizing a specific treatment protocol.

The present invention is directed to a method of inhibiting neoplastic cells by exposing those cells to a pharmacologically effective amount of pharmaceutical formulations comprising compositions containing those compounds of the present invention with pharmaceutical acceptable additives, diluents, carriers and excipients and at least one chemotherapeutic agent. Such compounds are effective at eliminating and inhibiting the growth of neoplasias such as precancerous lesions, tumours and cancer growth. One of the advantages of utilising such compositions is that they are low in toxicity, which in combination with their mechanism of action diminishes resistance development.

The present invention provides the compounds of the present invention, circiliol, including their precursor molecules, derivatives, metabolites, analogues and/or mimic molecules with pharmaceutical acceptable additives, diluents, carriers and excipients and pharmaceutically acceptable salts thereof, as well as pharmaceutical compositions comprising the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers and excipients and at least one chemotherapeutic agent and methods comprising inhibiting tumour growth or treating cancer by administering one or more of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers and excipients in combination with at least one chemotherapeutic agent.

The present invention also provides products that are useful for treating neoplasia and/or viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation with minimal toxic side effects unlike the high toxicity associated with standard chemotherapeutic agents.

The present invention is also directed to a pharmaceutical composition comprising a biologically active amount of at least one compound of the present invention.

The present invention is also directed to providing pharmaceutical formulations for treating various neoplasias and/or viral, bacterial and/or parasite infections and/or suppression of immune response rejection in tissue transplantation.

5 The present invention is also directed to providing a method for treating various neoplasias and/or viral, bacterial and/or parasite infections and/or suppression of immune response rejection in tissue transplantation.

10 Treatment according to the present invention can be effected when the subject is a neonate. Administration is carried out prior to delivery of the neonate and/or during delivery of the neonate. Prophylactic treatment of HIV<sup>+</sup> pregnant mothers prevent the viral transmission to the fetus by being able to administer lower less toxic levels of chemotherapeutic agents in combination with the compounds of the present invention.

15 These and other objects of the present invention will become apparent from the description of the invention disclosed below, which descriptions are intended to limit either the spirit or scope of the invention but are only offered as illustrations of the preferred embodiments of the invention.

20 The present invention is directed to the treatment, inhibition and/or prevention of viral, bacterial and/or parasite infections and/or suppression of immune response rejection in tissue transplantation and/or tumours and/or cancer growth and more particularly to treating neoplasia and/or HIV.

25 The invention features a method of treating neoplasia and/or viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation, which includes administering to an animal an effective amount of a pharmaceutical formulation comprising a composition comprising at least one of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof, at least one chemotherapeutic agent and a pharmaceutically acceptable carrier.

30 The invention is directed to a method of treating tumours and/or viral, bacterial and/or parasite infections and/or suppression of immune response rejection in tissue transplantation comprising administering a biologically active amount of a pharmaceutical formula comprising a composition comprising at least one of the



compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof, at least one chemotherapeutic agent and a pharmaceutically acceptable carrier.

5 The invention features a method of treating cancer and/or viral, bacterial and/or parasite infections and/or suppression of immune response rejection in tissue transplantation comprising administering to a patient in need thereof a cancer and/or viral, bacterial and/or parasite treatment and/or suppression of immune response rejection in tissue transplantation an effective amount of a pharmaceutical formula comprising a composition comprising at least one of the  
10 compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof, at least one chemotherapeutic agent and a pharmaceutically acceptable carrier.

The inventions objects include the provision of pharmaceutical formulations and compounds, which are suitable for making the formulations. Other objects are to  
15 provide methods to use the formulations.

It was surprisingly found that when the compounds of the present invention were administered in combination with chemotherapeutic agents, the proliferation of neoplastic cells was inhibited, which is manifested, pursuant to one aspect of the present invention, in a broad-spectrum anti-neoplastic activity with minimal toxicity.

20 The compounds of the present invention are individually diverse, but collectively all act to inhibit the propagation of neoplastic cells, cancers, cancer growth and/or tumours.

The invention also features a method of treating neoplasia and/or viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue  
25 transplantation comprising administering to a patient an effective amount of a pharmaceutical composition including at least one compound of the present invention, at least one chemotherapeutic agent and a pharmaceutically acceptable carrier.

Treating neoplasia in a patient includes achieving, partially or substantially, one or  
30 more of the following: arresting the growth or spread of a cancer, reducing the extent of a cancer (e.g., reducing size of a tumour or reducing the number of

affected sites), inhibiting the growth rate of a cancer, and ameliorating or improving a clinical symptom or indicator associated with a cancer (such as tissue or serum components).

5 In accordance with the present invention, a method is provided to treat or prevent viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation comprising administering to a subject an effective amount of one or more of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers and excipients and pharmaceutically acceptable salts thereof in combination with a chemotherapeutic agent.

10 The present invention also provides the use of one or more of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers and excipients and pharmaceutically acceptable salts thereof in combination with a chemotherapeutic agent, for the manufacture of a medicament for a viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation.

15 The present invention also provides compounds for use in a method of treatment of a viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation, said method comprising administering one or more to a subject.

#### DETAILED DESCRIPTION OF THE INVENTION

25 The compounds of the present invention comprise circiliol and corresponding pharmaceutically acceptable precursor molecules, derivatives, metabolites, analogues, mimic molecules and mixtures thereof.

The present invention relates to a composition comprising one or more of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof.

30 In another embodiment of the invention, the composition includes but is not limited to one or more of the following components: Circiliol; 6,7-dimethoxy-5,3'4'-

trihydroxyflavone; 6-hydroxy-2,3,4-trimethoxy acetophenone; 2-(3,4-dibenzoyloxybenzoyloxy)-4,5,6-trimethoxy acetophenone; 3'4'-dibenzoyloxy-2-hydroxy-4,5,6-trimethoxydibenzoyl methane; 6,7-dibenzoyloxy-5,6,7-trimethoxy flavone; 3,4-dihydroxy-5,6,7-trimethoxy flavone; 3,4-diacetoxy-5,6,7-trimethoxy flavone, their  
5 derivatives, metabolites, analogues and/or mimic molecules with pharmaceutical acceptable additives, diluents, carriers and excipients and pharmaceutically acceptable salts thereof.

In one embodiment, the chemotherapeutic agents are antimetabolites, nucleotide and/or nucleoside analogues. Novel antimetabolites, nucleotide and/or nucleoside  
10 analogues are disclosed in US 6,127,540; US 5,837,871; US 5,879,700; US 5,977,061; US 5,981, 507; US 5,994,321; and US 6,071,922 and are incorporated by reference.

In another embodiment the nucleotide analogues are pyrimidine antagonists, preferably selected from but not limited to the group comprising: 5-flourouracil;  
15 cytosine arabinoside; Azacitidine. In another embodiment the nucleotide analogues are purine antagonists, preferably selected from but not limited to the group comprising: 6-mercaptopurine; azathioprine; 5-iodo-2'-deoxyuridine; 6-thioguanine; 2-deoxycoformycin, cladribine, cytarabine, fludarabine, mercaptopurine, thioguanine, pentostatin.

In another embodiment the nucleoside analogues are preferably selected from but not limited to the group comprising: purine and pyrimidine nucleosides; ara-nucleosides; amino-nucleosides; aza-nucleosides; In another embodiment the nucleoside analogues are preferably selected from but not limited to the group  
20 comprising: AZT (zidovudine); ACV; valacyclovir; famciclovir; acyclovir; cidofovir; penciclovir; ganciclovir; Ribavirin; ddC; ddl (zalcitabine); lamuvidine; Abacavir; Adefovir; Didanosine; gemcitabine; d4T (stavudine); 3TC; BW 1592; PMEA/bis-POM PMEA; ddT, HPMPG, HPMPA, PMEA, PMEG, dOTC; DAPD; Ara-AC, pentostatin; dihydro-5-azacytidine; tiazofurin; sangivamycin; Ara-A (vidarabine); 6-MMPR; 5-FUDR (floxuridine); cytarabine (Ara-C; cytosine  
25 arabinoside); 5-azacytidine (azacitidine); HBG [9-(4-hydroxybutyl)guanine],  
30 (1S,4R)-4-[2-amino-6-cyclopropyl-amino)-9H-purin-9-yl]-2-cyclopentene-1-me

thanol succinate ("159U89"), uridine; thymidine; idoxuridine; 3-deazauridine; cyclocytidine; dihydro-5-azacytidine; tricitabine, ribavirin and fludrabine.

In another embodiment, the chemotherapeutic agents selected from the group but are not limited to: Chloroquin, primaquine, mefloquine, pyrimethamine-sulfadoxone, atoraquone/dapsone; halofantrine; artemisinin derivatives; atoraquone + proguanil, co-artemether; podophyllotoxin; pentamidine, diloxanide furoate, metronidazole, tindazole, tetracycline, quinacrine, stibogluconate, amphotericin B, quinine, doxycycline, trimethoprim-sulfamethoxazole, metronidazole, nifurtimox, suramin, melarsoprol, benznidazole, their metabolites, salts derivatives or any other anti-parasitic agent thereof.

In another embodiment, the chemotherapeutic agents selected from the group but are not limited to: mitomycin C, nalidixic acid, puromycin, sanamycin, and actinomycin.

In another embodiment, the chemotherapeutic agents selected from the group but are not limited to: N<sup>6</sup>-( $\Delta^2$ -isopentyl) adenosine, N<sup>6</sup>-( $\Delta^2$ -isopentyl) adenosine-5'-monophosphate, N<sup>6</sup>-( $\Delta^2$ -isopentyl) adenosine-3',5'-cyclic monophosphate, benzyladenosine, N<sup>6</sup>-benzyladenosine-5'-monophosphate, N<sup>6</sup>-benzyladenosine-3',5'-cyclic monophosphate, furfuryladenosine, N<sup>6</sup>-furfuryladenosine-5'-monophosphate, N<sup>6</sup>-furfuryladenosine-3',5'-cyclic monophosphate, N-(purin-6ylcarbamoyl)-o-chloroaniline ribonucleoside, N-(purin-6ylcarbamoyl)-o-chloroaniline ribonucleoside-5'-monophosphate, N<sup>6</sup>-adamantyladenosine, N<sup>6</sup>-adamantyladenosine-5'-monophosphate, N-(purin-6ylcarbamoyl)-o-octylamine ribonucleoside, N-(purin-6ylcarbamoyl)-o-octylamine ribonucleoside-5'-monophosphate, N-(purin-6ylcarbamoyl)-o-octylamine ribonucleoside-3',5'-cyclic monophosphate, N<sup>6</sup>-( $\Delta^2$ -isopentyl)-2-methylthioadenosine, N<sup>6</sup>-(4-hydroxy-3-methyl-trans-2-butenyl)adenosine, N<sup>6</sup>-(3-chloro-trans-2-butenyl) adenosine, surfinal adenosine and preferred metabolites including N<sup>6</sup>-( $\Delta^2$ -isopentyl) adenine, 6-N-(3-methyl-3-hydroxybutylamino) purine, adenine, hypoxanthine, uric acid and methylated xanthines.

In a further or alternative aspect of the invention, there is provided a method for the treatment of a viral, bacterial and/or parasite infection and/or suppression of immune response rejection in tissue transplantation comprising the step of administering an effective dose of a compound of the present invention as defined hereinabove or a

pharmaceutically acceptable derivative thereof combined with at least one  
chemotherapeutic agent. By combining the compounds of the present invention with  
chemotherapeutic agents a synergistic effect is observed and a much lower dosage  
of the chemotherapeutic agents is required, achieving the desired therapeutic effect  
5 while limiting the toxicity to the patient.

As will be appreciated by those skilled in the art, references herein to treatment  
extends to prophylaxis as well as to the treatment of established infections of  
symptoms.

10

By the term "pharmaceutically acceptable derivative" is meant any pharmaceutically  
acceptable salt, ester, or salt of such ester of a compound of the present invention, or  
any other compound which, upon administration to the recipient, is capable of  
providing (directly or indirectly) a compound of the present invention or an active  
15 metabolite or residue thereof.

Pharmaceutically acceptable salts of the compounds of the present invention include  
those derived from pharmaceutically acceptable inorganic and organic acids and  
bases. Examples of suitable acids include hydrochloric, hydrobromic, sulphuric, nitric,  
20 perchloric, fumaric, maleic, phosphoric, lactic, salicylic, succinic, p-toluenesulphonic,  
tartaric, acetic, citric, methanesulphonic, formic, benzoic, malonic, naphthalene-2-  
sulphonic and benzenesulphonic acids. Other acids such as oxalic, while not in  
themselves pharmaceutically acceptable, may be useful in the preparation of salts  
useful as intermediates in obtaining the compounds of the invention and their  
25 pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g. sodium), alkaline earth  
metal (e.g. magnesium), ammonium and  $N(R')_{4+}$  (where  $R'$  is  $C_{1-4}$   
alkyl) salts.

30 In a preferred embodiment, the nucleoside analogue is a phosphate ester selected  
from the group comprising: Acyclovir; 1-beta-D-arabinofuranosyl-E-5-(2-  
bromovinyl)uracil; 2'-fluorocarbocyclic-2'-deoxyguanosine; 6'-fluorocarbocyclic-2'-  
deoxyguanosine; 1-(beta-D-arabinofuranosyl)-5(E)-(2-iodovinyl)uracil; {(1r-1.alpha.,

2.beta., 3.alpha.)-2-amino-9-(2,3-bis(hydroxymethyl)cyclobutyl)-6H-purin-6-one}  
Lobucavir; 9H-purin-2-amine, 9-((2-(1-methylethoxy)-1-((1-  
methylethoxy)methyl)ethoxy)methyl)-(9Cl); trifluorothymidine; 9- $\rightarrow$ (1,3-dihydroxy-2-  
propoxy)methylguanine (ganciclovir); 5-ethyl-2'-deoxyuridine; E-5-(2-bromovinyl)-  
5 2'-deoxyuridine; 5-(2-chloroethyl)-2'-deoxyuridine; buciclovir; 6-deoxyacyclovir; 9-(4-  
hydroxy-3-hydroxymethylbut-1-yl)guanine; E-5-(2-iodovinyl)-2'-deoxyuridine; 5-  
vinyl-1-beta-D-arabinofuranosyluracil; 1-beta-D-arabinofuranosylthymine; 2'-nor-  
2'deoxyguanosine; 1-beta-D-arabinofuranosyladenine.

In another embodiment, the nucleoside phosphate ester is in the form of a  
10 pharmaceutically acceptable salt. In another embodiment, the salt is selected from  
the group consisting of sodium, potassium, ammonium, and hydrogen salts.

In certain embodiments the compounds of the present invention and  
chemotherapeutic agents are combined with a physiological carrier for the treatment  
15 of the pathophysiological state.

In another embodiment the pharmaceutical formulation is delivered to infected cells  
by incorporating the pharmaceutical formulation into liposomes or carbohydrate  
vehicles.

In another embodiment the pharmaceutical formulation further includes a  
pharmaceutically acceptable carrier, which in one embodiment is a cyclodextrin,  
alpha-cyclodextrin, beta-cyclodextrin, (beta-hydroxypropylcyclodextrin) gamma-  
cyclodextrin and in another embodiment is vitamin E oil, DMSO and/or ethanol.

25 The compounds of the present invention can be formulated and administered as  
free bases or in the form of their pharmaceutically acceptable salts for purposes of  
stability, convenience of crystallisation, increased solubility, and the like.

As used herein, the term "neoplasia" or neoplasm covers dysplasia, precancerous  
lesions, cancerous lesions, neoplastic cells, cancer, cancer growth, tumours,  
30 benign tumours, malignant tumours, solid tumours, carcinomas, etc.

As used herein, the term "precancerous lesion" includes syndromes represented by  
abnormal neoplastic, including dysplastic, changes of tissue. Examples include

precancerous growths in colonic, breast, renal, central nervous, gastric, or lung tissues, or conditions such as dysplastic nevus syndrome, a precursor to malignant melanoma of the skin. Examples also include, in addition to dysplastic nevus syndromes, polyposis syndromes, colonic polyps, precancerous lesions of the cervix (i.e., cervical dysplasia), prostatic dysplasia, bronchial dysplasia, breast, bladder and/or skin and related conditions (e.g., actinic keratosis), whether the lesions are clinically identifiable or not.

The pharmaceutical formulations of the present invention can be administered to a mammal having a susceptible cancer, i.e. a malignant cell population or tumour.

10 The combination of the compounds of the present invention and chemotherapeutic agents are effective on human tumours in vivo as well as on tumour cell lines in vitro. The pharmaceutical formulations of the present invention may be particularly useful for the treatment of solid tumours for which relatively few treatments are available. Such tumours include epidermoid and myeloid tumours, acute or

15 chronic, nonsmall cell, squamous. Specific cancers which may be mentioned as susceptible to treatment by administration of pharmaceutical formulations in accordance with the present invention include prostate cancer, colon cancer, small cell lung cancer, large cell lung cancer, lung adenocarcinoma, epidermoid lung cancer, melanoma (including amelanotic subtypes), renal cell carcinoma, gastric carcinoma, cancers of the central nervous system (based on the likelihood that the compounds will cross the blood cell barrier) including brain tumours, neuroblastomas, gastric carcinoma, breast cancer, ovarian cancer, testicular cancer, lymphoma and leukaemia, oesophageal cancer, stomach cancer, liver cancers, prostate cancer, cervical cancer, adrenal cancer, oral or mucosal cancer,

20 bladder cancer, pancreatic cancer, lymphoma, Hodgkins disease, sarcomas. Haematopoietic cell cancers such as B cell leukaemia/lymphomas, myelomas, T-cell leukemias/lymphomas, small cell leukemias/lymphomas, null cell, sezary, monocytic, myelomonocytic and Hairy cell leukemias. These lymphomas/leukemias can be either acute or chronic. Other cancers may also be

25 susceptible to treatment with the compounds of the present invention. The activity can readily be measured using standardised tests known to those skilled in the art.

30

As used herein, the term "carcinomas" refers to lesions that are cancerous. Examples include malignant melanomas, breast cancer, and colon cancer. As used herein, the term "neoplasm" refers to both precancerous and cancerous lesions.

5 As used herein, the terms "inhibit" or "inhibiting," mean decreasing tumour cell growth rate from the rate that would occur without treatment and/or causing tumour mass to decrease. Inhibiting also includes causing a complete regression of the tumour. Thus the compounds of the present invention can be either cytostatic or cytotoxic to the tumour cells.

10 In another embodiment, the subjects viral infection is selected from a DNA virus infection or an RNA virus. In a preferred embodiment, the DNA virus infection or the RNA virus infection is selected from a HIV, SHIV, SIV, FIV, HSV, CMV, HAV, HBV, HCV, HDV, HEV, EBV, BVDV, HSV-1, HSV-2, HSV-6, HHV-6, HHV-8, retrovirus infection, a togavirus infection, a flavivirus infection, a rubivirus infection,  
15 a pestivirus infection, a lipid envelope virus infection, a filovirus, a picornavirus infection, a rhinovirus infection, a coronavirus infection, a respiratory syncytial virus infection, a poliovirus infection, a parainfluenza virus infection, influenza virus infection, hantavirus, adeno-associated virus, measles virus, poxvirus, filovirus, human papilloma virus and animal papilloma virus infection.

20 In another embodiment, the patient is suffering from one or more complications or co-infections associated with AIDS, AIDS related syndromes, including cachexia and/or wasting syndrome.

25 In another embodiment, the parasite infection is a *Trypanosoma*, *Plasmodium*, *Cryptosporidium*, *Entamoeba*, *Balantidium*, *Leishmania*, *Pneumocystis*, *Trichomoniasis* or *Toxoplasma* parasite infection.

The present invention also relates to the administration of a pharmaceutical  
30 formulation to a patient infected with a retrovirus or is suffering with AIDS and co-infected with *Trypanosoma*, *Plasmodium*, *Cryptosporidium*, *Entamoeba*, *Balantidium*, *Leishmania*, *Pneumocystis*, *Trichomoniasis* or *Toxoplasma* parasites



or a *Mycoplasma* bacterium.

Another aspect of the present invention is the effective treatment of malaria, sleeping sickness, African trypanosomiasis, Chagas disease, American trypanosomiasis, cryptosporidiosis, amebiasis, balantidiasis, giardiasis, leishmaniasis, pneumocystosis, trichomoniasis, and toxoplasmosis with the pharmaceutical formulation of the present invention.

Accordingly, the present invention contemplates administering daily to a subject an amount of pharmaceutical formulation of the present invention that is clinically effective in treating or preventing a parasite infection, which the subject suffers or is at risk from infection. Illustrative parasites against which the invention can be applied are *Trypanosoma*, *Plasmodium*, *Cryptosporidium*, *Entamoeba*, *Balantidium*, *Leishmania*, *Pneumocystis*, *Trichomoniasis* or *Toxoplasma* parasites wherein the *Trypanosoma* parasites are selected from the group, but is not limited to *Trypanosoma cruzi*, *Trypanosoma brucei*, *Trypanosoma gambiense*, *Trypanosoma rhodesiense*, wherein the *Plasmodium* parasites are selected from the group, but is not limited to *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, *Plasmodium ovale*, *Plasmodium berghei*, wherein the *Entamoeba* is *Entamoeba histolytica*, wherein the *Balantidium* is *Balantidium coli*, wherein the *Leishmania* is selected for the group, but is not limited to *Leishmania braziliensis*, *Leishmania mexicana*, *Leishmania donovani*, *Leishmania tropica*, wherein the *Pneumocystis* is *Pneumocystis carinii*, wherein the *Trichomoniasis* is *Trichomoniasis vaginalis*, and the *Toxoplasma* parasite is *Toxoplasma gondii*, among other parasites.

In another embodiment, the bacterial infection is an intracellular bacterial infection, an extracellular bacterial infection, a mycoplasma infection, a *Listeria* infection or a *Mycobacterium* infection; a *Streptococcus* infection, a *Staphylococcus* infection, a *Vibrio* infection, a *Salmonella* infection; a *Shigella* infection, an enterotoxigenic, enteropathogenic, enteroinvasive or enterohemorrhagic *E. coli* infection, a *Yersinia* infection, a *Campylobacter* infection, a *Pseudomonas* infection, a *Borrelia* infection, a *Legionella* infection and a *Haemophilus* infection; pulmonary *Aspergillosis*, mucosal or oropharyngeal candidiasis and juvenile paracoccidiomycosis; or any

combination of the above.

In another embodiment, the composition further includes a pharmaceutically acceptable carrier.

5

In another embodiment, a pharmaceutical formulation or method in the prophylaxis and therapy of viral infections or a complication or consequence thereof is provided.

In particular the invention relates to the use a pharmaceutical formulation of the present invention, in the prophylaxis and therapy of viral infections, viral replication and the development and prevention of the deficiency of the immune system resulting in the development of opportunistic infections and certain cancers. More especially the invention relates to the use of a pharmaceutical formulation of the present invention, in the prophylaxis and therapy of viral infections, an example of which is the retrovirus, thought to be responsible for the Acquired Immune Deficiency Syndrome (AIDS) and AIDS related syndromes, believed to result from infection from the Human Immunodeficiency Virus (HIV), antibodies to which are found in almost all individuals diagnosed with AIDS.

Other embodiments provide a pharmaceutical formulation or method to treat a viral infection or to ameliorate one or more symptoms associated with a viral infection such as a flaviviral or retroviral infection comprising administering to an infected patient an effective amount of a pharmaceutical formulation as disclosed herein. It is an object of the present invention to provide a pharmaceutical formulation of the present invention for treatment, therapeutic or prophylaxis, against a viral, bacterial and/or parasitic infection. The compounds of the present invention are synergistic agents that enhance the efficacy of the chemotherapeutic agents, enabling their administration at much lower doses to achieve the therapeutic benefit, while minimizing the toxicity.

Accordingly disclosed herein is a pharmaceutical formulation or a method for treating a viral, parasite and/or bacterial infection and/or suppression of immune response rejection in tissue transplantation; for use in treating any infection, preventing a future infection and/or minimizing the effects of a future infection by a

virus, bacteria and/or parasite comprising administering to a patient in need thereof a prophylactically or therapeutically effective amount of a composition comprising at least one of the compounds of the present invention, with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof, at least one  
5 chemotherapeutic agent and a pharmaceutically acceptable carrier.

The advantage of this is that an effective anti-viral is provided that has minimal risk of conferring resistance and incurring toxicity.

10 In one embodiment of the invention the infection is a viral infection, in another embodiment the viral infection is caused by a retrovirus, in another embodiment, the retroviral infection is caused by HIV or AIDS virus, Herpes virus, cytomegalovirus, or an animal virus, and in an additional embodiment the viral infection is caused by a lipid envelope virus.

15 In another embodiment, the pharmaceutical formulation is used to treat AIDS related syndromes, including cachexia and/or wasting syndrome.

20 One useful property of the pharmaceutical formulation of the present invention is their anti-viral activity, which is manifested, pursuant to one aspect of the present invention, in a broad-spectrum anti-viral activity. Accordingly, the present invention contemplates administering daily to a subject an amount of a pharmaceutical formulation of the present invention that is clinically effective in treating or preventing a viral infection, which the subject suffers or is at risk from infection.

25 Illustrative viruses against which the invention can be applied are HIV, cytomegalovirus (CMV), a KS-producing herpes virus, Kaposi's Sarcoma-associated herpes virus, the virus of the genus hepatitis, a virus of the genus picornaviruses, a virus of the genus molluscipoxvirus, hantaviruses, among other viruses.

30 Additionally, the present invention is also broadly directed providing the use of the pharmaceutical formulation of the present invention in the treatment (i.e., in the sense of treating an existing infection, preventing a future infection, minimizing the

effect of a future infection and/or enhancing the performance of a vaccine) of all infections which are not retroviral infection, several representative examples of which include one or more kind of Mycoplasma, and/or one or more diseases caused by Mycoplasmas and/or one or more of the following infections: hairy  
5 leukoplakia, oral candidosis, mouth ulcerations – aphthous/herpetic/bacterial, fungal candida, squamous oral carcinoma, Kaposi's sarcoma oral lesions, periodontitis, necrotizing gingivitis, human papilloma virus, rhinovirus and arboviral molluscum contagiosum, orafacial herpes zoster, Epstein barr virus, rotaviruses, togaviruses, including alpha viruses (also known as arboviruses, group A),  
10 flaviviruses (also known as arboviruses, group B, such as yellow fever, as well as hepatitis C and hepatitis G), rubiviruses (also known as rubella viruses e.g., human rubella virus), pestiviruses (also known as mucosal disease viruses, such as bovine virus diarrhoea virus BVDV, hog cholera virus, and sheep border disease), as well as any other non-retroviral viral induced infections. Thus virus infections that may  
15 be treated include but are not limited to HIV, SIV, FIV, FELV, SHIV, Kaposi's Sarcom-associated herpes virus and other herpes viruses (e.g. HSV-1, HSV-2, human herpes virus 6 (HHV-6) and HHV-8), the viruses associated with hepatitis (HAV, HBV, hepatitis C virus [HCV]), and human cytomegalovirus, togaviruses and flaviviruses, e.g., California encephalitis virus, St. Louis encephalitis virus, western  
20 equine encephalitis virus, eastern equine encephalitis virus, Colorado tick fever virus, LaCrosse encephalitis virus, Japanese encephalitis virus, yellow fever virus, Venezuelan equine encephalitis virus, Murray valley fever virus, tick-borne encephalitis virus, GB virus A, GB virus B, GB virus C, Dengue virus 1, Dengue virus 2, Dengue virus 3, Dengue virus 4, Semliki Forest virus, Sinbis virus,  
25 picornaviruses, rhinoviruses, coronaviruses, respiratory syncytial viruses, polioviruses, parainfluenza viruses and influenza viruses ( including type A, type B, and type C.

Pursuant to a preferred embodiment of the invention, an effective amount of a  
30 pharmaceutical formulation of the present invention thus administered is such as to produce a circulating concentration of the compounds of the present invention, including precursor molecules, derivatives, metabolites, analogues, mimic molecules,

and at least one chemotherapeutic agent sufficient to reduce viral loads as monitored by, e.g., viral titre methods or by PCR.

According to a further aspect of the invention the pharmaceutical formulations is formulated in a liposome.

5

In another embodiment of the invention, liposomes are provided carrying the pharmaceutical formulations of the present invention targeted to HIV infected cells by putting antibodies to the HIV coat protein gp160 or gp120 on its surface. The advantage of this is that the liposome can selectively target HIV infected cells.

10 As used herein, the terms subject and patient are used interchangeably. Subjects and patients are mammals.

The pharmaceutical formulations of the present invention are useful antineoplastic agents i.e. to inhibit tumour cell growth in vitro and in vivo, in mammalian hosts, such as humans or domestic animals, and are particularly effective against solid  
15 tumours and multidrug resistant tumours. Thus the invention provides a method comprising inhibiting cancer cells, by contacting said cells, in vitro and in vivo with an effective amount of a pharmaceutical formulation of the present invention. The invention also provides a therapeutic method comprising treating cancer (i.e. inhibiting tumour cell growth) by administering a pharmaceutical formulation of the  
20 present invention to a mammal in need of such therapy.

The present invention is also directed to pro-drug compounds analogous to the active compounds disclosed herein. Such compounds are generally themselves inactive or low in activity, but are converted into active compounds. Thus, for example, pro-drugs such as the methyl ester of any acid functionality, which is not  
25 active *per se* or has very low activity could be hydrolysed, either uncatalytically or catalytically with an enzyme such as an esterase, to an active compound. Such pro-drug compounds could well be the preferred therapeutic form of the present compounds. These analogous prodrugs can be produced from active compounds based on procedures and factors that are well known to one of ordinary skill in the  
30 art. Accordingly as used in the present application, "pro-drug analogue" means "a chemical which is relatively non-toxic and pharmacologically inert but which can be transformed *in vivo* to a pharmacologically active drug". More specifically it means

a precursor molecule, derivative, metabolite or analogue of the compounds of the present invention which have low or no ability as antineoplastic, antiviral, antibacterial, antiparasitic agents until converted in the body to a derivative, metabolite or analogue with such ability or abilities. Such pro-drugs should have  
5 favourable properties such as enhanced absorption, water solubility, lower toxicity, or better targeting to the tumour cell (such as by reason of greater affinity to the tumour cell or a larger quantity of activating enzyme in the tumour cell as opposed to a normal cell so that larger concentrations of the active compound are produced in the tumour cell). Examples of such compounds are esters, such as methyl,  
10 ethyl, phenyl, N,N-dimethylaminoethyl, acyl derivatives such as benzoyl, p-N,N-dimethylaminobenzoyl, N,N-dimethylaminoglycyl, peptide derivatives such as  $\gamma$ -glutamyl, glycyl, D-Val-Leu-Lys.

The pharmaceutical formulation according to the present invention can be administered to a patient in any of a wide range of routes. Thus, with regard to the  
15 types of formulations in which the active compounds according to the present invention can be administered, as well as any additives can be included with the active compounds in the formulations, and the possible routes of administration, it is well known to those of skill in the art that such formulations can be provided in a wide variety of types, and it is within the skill of the ordinary artisans to select a  
20 specific formulation and route of administration and then test suitability for use. By way of example but not limitation, suitable routes include enteric, parenteral, topical, oral, rectal, nasal or vaginal routes. Parenteral routes include subcutaneous, intramuscular, intravenous, intraperitoneal, intradermal and sublingual administration. Also, compositions may be implanted into a patient or  
25 injected using a drug delivery system.

The pharmaceutical formulation according to the present invention may be administered locally or systemically. By systemic administration means any mode or route of administration that results in effective amounts of active ingredient appearing  
30 in the blood or at a site remote from the route of administration of the active ingredient.

Further, the pharmaceutical formulation according to the present invention may be

administered intermittently. The advantage of this is that it allows the patient to suspend therapy for periods without the worry of inactivity of the drug resulting from the development of resistant cells.

- 5     The pharmaceutical formulation according to the invention may be formulated for enteral, parenteral or topical administration. Indeed all three types of formulations may be used simultaneously to achieve systemic administration of the active ingredient.

10     Compounds useful in the methods of this invention may be formulated into compositions together with pharmaceutically acceptable carriers for oral administration in solid or liquid form, or for rectal administration, although carriers for oral administration are most preferred.

15     Pharmaceutically acceptable carriers for rectal administration are preferably suppositories that may contain, in addition to the compounds of the present invention, excipients such as cocoa butter or a suppository wax.

20     Pharmaceutically acceptable carriers for oral administration include capsules, tablets, pills, powders, troches and granules. In such solid dosage forms, the carrier can comprise at least one inert diluent such as sucrose, lactose or starch. Such carriers can also comprise, as is normal practice, additional substances other than diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, troches and pills, the carriers may also comprise buffering agents. Carriers such as tablets, pills and granules can be prepared with enteric coatings on the surfaces of the tablets, pills or granules. Alternatively, the enterically coated compound can be pressed into a tablet, pill, or granule, and the  
25     tablet, pill or granules for administration to the patient. Preferred enteric coatings include those that dissolve or disintegrate at colonic pH such as shellac or Eudraget S. Additional pharmaceutically acceptable carriers include liquid dosage forms for oral administration, e.g. pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs containing inert diluents commonly used  
30     in the art, such as water. Besides such inert diluents, compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, and sweetening, flavouring and perfuming agents.

In one embodiment, the pharmaceutical formulation has an enteric coating. In another embodiment, the enteric coating is made of a polymer or copolymer. In a preferred embodiment, the polymer or copolymer is selected from the group consisting of poly(lactic-glycolic acid) polyester, cellulose acetate phthalate, hydroxypropyl-methyl cellulose phthalate poly(butyl methacrylate), (2-dimethyl aminoethyl) methacrylate, and methyl methacrylate.

In accordance with the present invention, the expression "micronized" means that the compound has been micronized in accordance with any process for micronizing, a number of which are known in the art. The micronized particles preferably include a percentage of particles, which are of a diameter, which is about 10 microns, or less, preferably, 5 microns or less. For example, in a preferred aspect of the invention, at least 80% of the particles in a formulation of micronized particles have a diameter of less than 5 microns. An alternative to micronizing a compound is to solubilize the compound and put it into liposomes of appropriate size. The manufacture of liposomes and the insertion of active ingredients into such liposomes are well known in the art.

Suitable injectable solutions include intravenous, subcutaneous and intramuscular injectable solutions. Examples of injectable forms include solutions, suspensions and emulsions. Typically the compound(s) is injected in association with a pharmaceutical carrier such as normal saline, Ringers solution, dextrose solution and other aqueous carriers known in the art. Appropriate non-aqueous carriers may also be used and examples include cyclodextrin, preferably hydroxypropyl beta cyclodextrin, mixed oils (vitamin E oil), polyethylene glycol and ethyl oleate. A preferred carrier is cyclodextrin in water. Frequently, it is desirable to include additives in the carrier such as buffers and preservatives or other substances to enhance isotonicity and chemical stability.

The pharmaceutical formulation can also be administered topically. Suitable formulations for topical administration include creams, gels, jellies, muclriages, pastes and ointments. The compounds may be formulated for transdermal administration, for example in the form of transdermal patches so as to achieve systemic administration.

The pharmaceutical formulation may also be administered in the form of an implant.



The pharmaceutical formulation may also be administered in the form of an infusion solution or as a nasal inhalation, aerosol or spray.

5 In another embodiment, the composition is incorporated in a pharmaceutically acceptable carrier, diluents, vehicles and the like for systemic administration by feeding. An example of such a carrier is cyclodextrin ( $\alpha$ -cyclodextrin,  $\beta$ -hydroxypropylcyclodextrin or  $\gamma$ -cyclodextrin), DMSO, ethanol.

10 The pharmaceutically acceptable carrier, chemotherapeutic agents and compounds of this invention are formulated into unit dosage forms for administration to a patient. The dosage levels of active ingredient (i.e. compounds of this invention) in the unit dosage may be varied so as to obtain an amount of active ingredient effective to achieve lesion-eliminating activity in accordance with the desired method of administration (i.e., oral or rectal). The selected dosage level therefore depends upon the nature of the active compound administered, the route of  
15 administration, the desired duration of treatment, individual needs and other factors. If desired, the unit dosage may be such that the daily requirement for active compound is in one dose, or divided among multiple doses for administration, e.g., two to four times per day.

20 With regard to dosage and duration of treatment according to any aspect of the present invention, it is recognized that the ability of an artisan skilled in pharmaceutical administration of drugs to determine suitable dosages depending on many inter-related factors is well known, and skilled artisans are readily able to monitor patients to determine whether treatment should be started, continued, discontinued or resumed at any given time. For example, dosages of the  
25 compounds are suitably determined depending on the individual cases taking symptoms, age and sex of the subject and the like into consideration. The amount of the compound to be incorporated into the pharmaceutical composition of the invention varies with dosage route, solubility of the compound, administration route, administration scheme and the like. An effective amount for a particular patient  
30 may vary depending on factors such as the condition being treated, the overall health of the patient and the method, route and dose of administration. The clinician using parameters known in the art makes determination of the appropriate

dose. Generally, the dose begins with an amount somewhat less than the optimum dose and it is increased by small increments thereafter until the desired or optimum effect is achieved. Suitable dosages can be determined by further taking into account relevant disclosure in the known art. In one embodiment, the unit dose  
5 comprises 5-500 mg of active ingredient consisting of at least one compound of the present invention.

The pharmaceutical formulation of this invention is preferably packaged in a container (e.g. a box or bottle, or both) with suitable printed material (e.g. a package insert) containing indications, directions for use, etc.  
10 The pharmaceutical formulations containing the active compounds or pro-drugs of the present invention can be formulated so as to be specifically targeted to tumours and viral, bacterial and/or parasite containing cells. The compounds can be attached to the reagent that is capable of binding a tumour and/or viral, bacterial and/or parasite-associated antigen. For example, the compounds of the present  
15 invention could be covalently attached to a monoclonal antibody such as directed to a tumour-associated antigen. The antigen may be located on a tumour/virus/ bacterial/parasite or in the tumour/virus cell area. Such linkages can be made through peptide bond formation with amino groups of an antibody. Suitable reagents include polyclonal and monoclonal antibodies. Accordingly, the present  
20 invention also provides a method comprising treating cancer and/or viral/ bacterial/parasite infections by administering a pharmaceutical formulation of the present invention and a reagent (i.e. monoclonal or polyclonal antibody), which is capable of binding to a tumour/virus/ bacterial/parasite associated antigen.

In one embodiment, the liposomes or carbohydrate vehicles are targeted to HIV  
25 infected cells by putting viral antibodies on its surface. In another embodiment, the viral antibodies are directed to the HIV coat protein gp160 and/or gp120.

The invention provides a method of treating a neoplasia and/or viral/ bacterial/parasite infection and/or suppression of immune response rejection in tissue transplantation comprising administering to a mammal in need thereof an  
30 effective amount of a composition containing as an active ingredient therein at least one of the compounds of the present invention, with pharmaceutical acceptable

additives, diluents, carriers, excipients and pharmaceutical salts thereof, a chemotherapeutic agent and a pharmaceutically acceptable carrier.

The invention provides a method for manufacturing a medicament for treating neoplasia and/or viral/ bacterial/parasite infection and/or suppression of immune response rejection in tissue transplantation wherein the treatment protocol comprises:

Administering a first composition comprising at least one of the compounds of the present invention circiliol, precursor molecules, derivatives, metabolites, analogues and/or mimic molecules with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof and a pharmaceutically acceptable carrier;

later followed by a second composition comprising at least one conventional chemotherapeutic agent and a pharmaceutically acceptable carrier;

during the treatment protocol.

The present invention also provides a method of endowing a chemotherapeutic agent with substantially enhanced therapeutic efficacy and reduced toxicity, comprising:

- (a) providing a chemotherapeutic agent; and
- (b) combining the agent with one or more compounds of the present invention with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof and a pharmaceutically acceptable carrier thereof so as to reduce the cytotoxicity of the combination in comparison to the chemotherapeutic agent alone.

The invention further provides a pharmaceutical formulation for treating a cancer selected from the group consisting of small cell lung cancer, testicular cancer,

lymphoma, leukaemia, oesophageal cancer, stomach cancer, colon cancer, breast cancer, central nervous system cancer, liver cancer and prostate cancer, which comprising administering to a mammal in need thereof an effective amount of a composition containing as an active ingredient therein at least one of the  
5 compounds of the present invention, their derivatives, metabolites, analogues and/or mimic molecules with pharmaceutical acceptable additives, diluents, carriers, excipients and pharmaceutical salts thereof.

The invention provides compositions comprising, the compounds of the present invention and at least one chemotherapeutic agent, to be used as anti-viral/  
10 bacterial/parasite and/or anti-cancer agents.

The invention provides compositions comprising, the compounds of the present invention and at least one chemotherapeutic agent, to be used in the preparation of medicaments having anti-viral/ bacterial/parasite and/or anti-cancer activity.  
15

The invention further provides, the use of the compounds of the present invention and at least one chemotherapeutic agent, is provided as anti-viral/ bacterial/parasite and/or anti-cancer agents.

20 In another embodiment, the use of the compounds of the present invention and at least one chemotherapeutic agent, is provided for the preparation of medicaments having anti-viral/ bacterial/parasite and/or anti-cancer activity.

In another embodiment, the compounds of the present invention and at least one chemotherapeutic agent, are provided, for the preparation of medicaments having activity against viral/ bacterial/parasite infections and/or cancer.  
25

The invention also relates to a method of suppressing tumour growth in a mammal by administering to the mammal an amount of the compounds of the present invention, and a second chemotherapeutic agent effective to suppress tumour growth in the mammal. The second chemotherapeutic agent is an antimetabolite, nucleotide and/or nucleoside analogue. These compositions provide enhanced antitumour effect and may also prevent the development of metastases. In  
30

particular, these compounds are useful for overcoming tumours that are drug resistant. These agents may be administered separately or as a cocktail. Administering the compound of the present invention or a derivative, metabolite, analogue or mimic molecule, thereof several hours prior to administering the chemotherapeutic agent may reduce toxicity. The compositions can be administered by any route.

The components of any of the pharmaceutical formulations disclosed herein can be administered simultaneously (in a combination formulation), essentially simultaneously (e.g., administration of each compound a few minutes or a few hours apart), or can be administered sequentially, e.g., several days apart, or more than a week apart. For example, a compound of the present invention, (and a conventional chemotherapeutic agent) can be administered together, or essentially simultaneously, e.g., administration of each compound a few minutes or a few hours apart, or can be administered sequentially, e.g., several days apart, or more than a week apart. All such variations in administration of the combination therapy are encompassed within the scope of the invention.

Further, the invention provides use of the composition in veterinary medicine, prophylactically and therapeutically in animal populations that are subject to infection that compromises immune response and cause infection.

In another embodiment, the subject or patient is an animal. The term animal as used herein includes, but is not limited to, mice, rats, domesticated animals such as but is not limited to, cats, dogs, and other animals but is not limited to, cattle, sheep, pigs, horses, camels, buffalo and primates such as but not limited to, monkeys, humans and more generally mammals.

Treatment according to the present invention can be effected when the subject or mammal is a neonate. Administration is carried out prior to delivery of the neonate and/or during delivery of the neonate.

Without further elaboration, it is believed that one skilled in the art can, using the

preceding description, practice the present invention to its fullest extent. The following detailed examples describe how to test the various compounds of this invention and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way  
5 whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures.

These and other features and advantages of the present invention will be more clearly understood with reference to the following description of some embodiments  
10 thereof and with reference to the accompanying drawings in which: -

Fig. 1 A and B are graphs showing the IC<sub>50</sub> and IC<sub>70</sub> values of gemcitabine alone and in combination with circiliol respectively; and

15 Fig. 2 is graphs showing the dose dependent influence of circiliol on gemcitabine antitumour activity in both cell lines.

#### EXAMPLE 1

Gemcitabine (2', 2'-difluoro-2'-deoxycytidine, dFdC) is approved for the treatment  
20 of locally advanced or metastatic pancreatic cancer, with additional activity in non small lung cancer alone and in combination with other cytotoxic agents like 5-flourouracil or cisplatin.

For this study, the *in vitro* growth of gemcitabine and circiliol, alone and in  
25 combination was studied in the large cell lung cancer line LXFL 529L and the lung adenocarcinoma cell line LXFA 526L in a monolayer proliferation inhibition study.

#### TEST PROCEDURE

##### Cell lines

30 The cells were grown in monolayers and routinely passaged weekly. Stocks were not maintained more than 20 passages *in vitro*. Table 1 summarises the cell line characteristics.

**TABLE 1: Characterisation of cell lines**

<b>Tumour type</b>	<b>Cell line</b>	<b>Histology in nude mice</b>	<b>Doubling time (hours)</b>	<b>Tumour formation in vivo</b>
Lung (NSC)	LXFA 526L	adenocarcinoma	34	yes
	LXFA 529L	large cell	25	yes

NSC = non small cell

### Assay

5 A modified propidium iodide assay (Dengler et al, 1995) was used to assess the effects of the 2 compounds and the combinations thereof. Cells were harvested from exponential phase cultures growing in RPMI 1640 medium supplemented with 10% foetal calf serum (FCS) and 1% gentamycin (100 U/ml) by trypsination, counted and plated in 96 well flat-bottomed microtitre plates (140  $\mu$ l cell  
10 suspension, 5-20,000 cells/well). After a 24 hour recovery, to allow the cells resume exponential growth, 10  $\mu$ l of culture medium (6 control wells per plate) or culture medium containing gemcitabine and/or circiliol were added to the wells. Each drug concentration was plated in triplicate. Following 4 days of continuous drug exposure, the culture medium with or without drug was removed and 200  $\mu$ l of  
15 an aqueous propidium iodide solution (6  $\mu$ g/ml) was added which then intercalates into the DNA of the dead cells. Fluorescence was measured using a Cytofluor 2350 microplate reader (excitation 530 nm, emission 620 nm). Growth inhibition was expressed as treated/control X 100 (%T/C); IC<sub>50</sub> and IC<sub>70</sub> values were determined by plotting drug concentration versus cell viability.

20

### Drug application.

Circiliol, a crystalline compound, was dissolved in DMSO at a stock concentration of 100 mg/ml and stored at 4°C. Gemcitabine was used as its clinical formulation (Gemzar, Lilly) and dissolved for each experiment freshly in sterile water at a stock  
25 concentration of 3.3 mg/ml.

### Results and discussion

Both cell lines grew very well, with initial cell number increasing 4 fold.

Fluorescence units of controls ranged from 1.421 to 4.277. Clear dose response effects were seen. The determination of antiproliferation effects were repeated up to five times. The *in vitro* activity of circiliol is summarized in table 2.

5 **TABLE 2: *In vitro* activity (Test/Control %) of Circiliol**

Concentration of Circiliol ( $\mu\text{g/ml}$ )	3	30	300
LXFA 526L	93	53	56
LXFA 529L	102	49	39

**MEDIAN OF 5 EXPERIMENTS**

10 Circiliol alone was only moderately active *in vitro*. Concentration of 30  $\mu\text{g/ml}$  and 300  $\mu\text{g/ml}$  resulted in tumour growth inhibition in the range of 39% to 56% versus control in both cell lines. At concentrations of  $\leq 3$   $\mu\text{g/ml}$  circiliol no antitumour activity was evident *in vitro*.

15 Gemcitabine was potently active in both cell lines *in vitro*. The mean  $\text{IC}_{50}$  value was 0.63  $\mu\text{g/ml}$  in LXFA 526L adenocarcinoma cell lines and 0.36  $\mu\text{g/ml}$  in the LXFL 529L large cell lung cancer cell lines. The mean  $\text{IC}_{70}$  value was 5.7  $\mu\text{g/ml}$  in LXFA 526L and 0.64  $\mu\text{g/ml}$  in LXFL 529L.

20 Based on these results, gemcitabine was combined with 3  $\mu\text{g/ml}$  circiliol, which showed no tumour effects, and 30  $\mu\text{g/ml}$  circiliol which resulted in 50% inhibition of tumour growth *in vitro*. The results are presented in tables 3 and 4 and figures 1 and 2.

**TABLE 3:  $\text{IC}_{50}$  values  $\mu\text{g/ml}$  of gemcitabine alone and in combination with Circiliol**

Concentration of Circiliol	0 $\mu\text{g/ml}$	+ 3 $\mu\text{g/ml}$	+ 30 $\mu\text{g/ml}$
LXFA 526L	0.63	0.072	n.d.
LXFA 529L	0.355	0.039	n.d.

25  **$\text{IC}_{50}$  VALUES EVALUATED AS A MEDIAN OF 2-5 EXPERIMENTS PER CELL**



**LINE.**

**n.d. NOT DETECTABLE**

**TABLE 4: IC<sub>70</sub> values  $\mu\text{g/ml}$  of gemcitabine alone and in combination with Circiliol**

Concentration of Circiliol	0 $\mu\text{g/ml}$	+ 3 $\mu\text{g/ml}$	+ 30 $\mu\text{g/ml}$
LXFA 526L	5.6	0.87	0.58
LXFA 529L	0.64	0.076	0.033

**IC<sub>70</sub> VALUES EVALUATED AS A MEDIAN OF 2-5 EXPERIMENTS PER CELL**

**LINE.**

Treatment of cells with 3  $\mu\text{g/ml}$  circiliol in combination with gemcitabine shifted IC<sub>50</sub> values from 0.63  $\mu\text{g/ml}$  to 0.072  $\mu\text{g/ml}$  in LXFA 526L cell lines and from 0.36  $\mu\text{g/ml}$  to 0.04  $\mu\text{g/ml}$  in the LXFA 529L cell lines. IC<sub>70</sub> values from 5.6  $\mu\text{g/ml}$  to 0.87  $\mu\text{g/ml}$  in LXFA 526L cell lines and from 0.64  $\mu\text{g/ml}$  to 0.076  $\mu\text{g/ml}$  in the LXFA 529L cell lines. Treatment of cells with 30  $\mu\text{g/ml}$  circiliol in combination with gemcitabine shifted IC<sub>70</sub> values from 5.6  $\mu\text{g/ml}$  to 0.58  $\mu\text{g/ml}$  in LXFA 526L cell lines and from 0.64  $\mu\text{g/ml}$  to 0.0033  $\mu\text{g/ml}$  in the LXFA 529L cell lines, corresponding to a 10 and 19 fold increase of antitumour activity respectively.

Circiliol alone demonstrates low activity in no small lung tumour cells LXFA 526L and LXFL 529L. However, when administered in combination with the antimetabolite gemcitabine a 6 to 8 fold increase in antitumour activity of gemcitabine is observed in vitro at a concentration of 3  $\mu\text{g/ml}$ , which is inactive alone.

Pharmaceutically acceptable refers to those properties and/or substances, which are acceptable to the patient from a pharmacological/toxicological point of view including bioavailability and patient acceptance or to the manufacturing chemist from a physical-chemical point of view regarding composition, formulation, stability and isolatability.

The terms "comprise, comprised and comprising" and the terms "include, included and including" are used interchangeably in this specification and are to be afforded the widest interpretation.

- 5 The invention is not limited to the embodiments described above, but may be varied in both construction and detail within the scope of the claims.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
222